

Exercises: Gradient algorithms

Exercise 1 (A quadratic example in \mathbb{R}^2). Consider, for $\gamma > 0$, $f(x) = \frac{1}{2}(x_1^2 + \gamma x_2^2)$. We apply the gradient descent method with optimal step, starting at $x^{(0)} = (\gamma, 1)$.

1. Show that f is m -convex with M -Lipschitz gradient. Find the tightest m and M constants.

2. Show that

$$x^{(k)} = \left(\gamma \left(\frac{\gamma-1}{\gamma+1} \right)^k, \left(-\frac{\gamma-1}{\gamma+1} \right)^k \right)$$

and

$$f(x^{(k)}) = \frac{\gamma(\gamma+1)}{2} \left(\frac{\gamma-1}{\gamma+1} \right)^{2k} f(x^{(0)})$$

3. Show that, on this example, the convergence is exactly linear, that is $f(x^{(k)}) - v^\sharp$ is a geometric series. Give its reason. Compare with the theoretical bound.

4. When is this algorithm fast and slow ?

Exercise 2 (Strongly convex - optimal step). Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a m -convex \mathcal{C}^2 function. Define, for given $x^{(0)}$,

$$\tilde{f}_k : t \mapsto f(x^{(k)} - t \nabla f(x^{(k)}))$$

$$t^{(k)} = \arg \min_{t \in \mathbb{R}} \tilde{f}_k(t)$$

$$x^{(k+1)} = x^{(k)} - t^{(k)} \nabla f(x^{(k)})$$

1. Show that there exists $M \geq m$ such that $mI \preceq \nabla^2 f(x^{(k)}) \preceq MI$

2. Show that, for any interesting t (to be defined) we have

$$\tilde{f}_k(t) \leq f(x^{(k)}) - t \|\nabla f(x^{(k)})\|_2^2 + \frac{Mt^2}{2} \|\nabla f(x^{(k)})\|_2^2$$

3. Show that,

$$f(x^{(k+1)}) \leq f(x^{(k)}) - \frac{1}{2M} \|\nabla f(x^{(k)})\|_2^2$$

4. Show that $f(x^\sharp) \geq f(x) - \frac{1}{2m} \|\nabla f(x)\|_2^2$

5. Show that

$$f(x^{(k+1)}) - f(x^\sharp) \leq (1 - \frac{m}{M}) [f(x^{(k)}) - f(x^\sharp)]$$

6. Show that the algorithm converges, and give its convergence speed.

Exercise 3 (Strictly convex case). Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a \mathcal{C}^2 , convex function with M -Lipschitz gradient such that $f(x^\sharp) = \inf f$. We define, for given $x^{(0)}$,

$$x^{(k+1)} = x^{(k)} - t \nabla f(x^{(k)})$$

with $t \leq 1/M$.

1. Show that, for all x and y $(y-x)^\top \nabla^2 f(z)(y-x) \leq M \|y-x\|_2^2$

2. Show that

$$f(y) \leq f(x) + \nabla f(x)^\top (y-x) + \frac{M}{2} \|y-x\|^2$$

3. Show that $f(x^{(k+1)}) \leq f(x^{(k)}) - \frac{t}{2} \|\nabla f(x^{(k)})\|^2$

4. Show that

$$f(x^{(k+1)}) \leq v^\sharp + \nabla f(x^{(k)})^\top (x^{(k)} - x^\sharp) - \frac{t}{2} \|\nabla f(x^{(k)})\|^2$$

5. Deduce that

$$f(x^{(k+1)}) \leq v^\sharp + \frac{1}{2t} (\|x^{(k)} - x^\sharp\|^2 - \|x^{(k+1)} - x^\sharp\|^2)$$

6. Show that

$$\sum_{i=1}^k f(x^{(i)}) - v^\sharp \leq \frac{1}{2t} \|x^{(0)} - x^\sharp\|^2$$

7. Conclude that

$$f(x^{(k)}) - v^\sharp \leq \frac{1}{2kt} \|x^{(0)} - x^\sharp\|^2.$$