

Exercises: Optimality conditions

February 23, 2026

Exercise 1. Solve the following optimization problem 2.

$$\begin{array}{ll}
 \text{Min}_{x,y \in \mathbb{R}^2} & (x-1)^2 + (y-2)^2 \\
 & x \leq y \\
 & x + 2y \leq 2
 \end{array}
 \qquad
 \begin{array}{ll}
 \min_{x_1, x_2, x_3} & 4x_1^2 - x_1x_2 + x_2^2 - 12x_1 \\
 \text{s.t.} & x_1 - 2x_2 + x_3 = 5 \\
 & x_1^2 + 3x_2^2 \leq 10 \\
 & x_1, x_2, x_3 \geq 0
 \end{array}$$

Exercise 2 (First order optimality condition). 3.
Consider, for f differentiable,

$$\begin{array}{ll}
 (P) \quad \text{Min}_{x \in \mathbb{R}^n} & f(x) \\
 \text{s.t.} & x \in X
 \end{array}
 \qquad
 \begin{array}{ll}
 \min_{x_1, x_2, x_3} & e^{x_1} - x_1x_2 + x_3^3 \\
 \text{s.t.} & \ln(e^{x_1-4x_2} + e^{x_1+x_3}) \leq 2x_1 + 3 \\
 & 2x_1^2 + x_2^2 \leq 2
 \end{array}$$

Recall that

$$T_X(x_0) = \left\{ d \in \mathbb{R}^n \mid \exists d_k \rightarrow d, \exists t_k \searrow 0, \text{s.t. } x_0 + t_k d_k \in X \right\}$$

and $K^\oplus = \{ \lambda \mid \lambda^\top x \geq 0, \forall x \in K \}$.

Show that

1. If x_0 is an optimal solution to (P), then $\nabla f(x_0) \in [T_X(x_0)]^\oplus$. 5.

2. If f is convex, X is closed convex, and $\nabla f(x_0) \in [T_X(x_0)]^\oplus$, then x_0 is an optimal solution to (P).

$$\begin{array}{ll}
 \min_{x_1, x_2} & -x_1 \\
 \text{s.t.} & -x_2 - (x_1 - 1)^3 \leq 0 \\
 & x_1, x_2 \geq 0
 \end{array}$$

$$\begin{array}{ll}
 \min_{x_1, x_2} & -x_1 \\
 \text{s.t.} & x_2 - (x_1 - 1)^3 \leq 0 \\
 & x_1, x_2 \geq 0
 \end{array}$$

Exercise 3. In the following cases, are the KKT conditions necessary / sufficient ?

1.

$$\begin{array}{ll}
 \min_{x_1, x_2, x_3} & 12x_1 - 5x_2 + 3x_3 \\
 \text{s.t.} & x_1 + 2x_2 - x_3 = 5 \\
 & x_1 - x_2 \geq -2 \\
 & 2x_1 - 4x_2 \leq 12
 \end{array}$$

Exercise 4. Solve the following problem using first order optimality conditions

$$\begin{array}{ll}
 \min_{x_1, x_2} & -2(x_1 - 2)^2 - x_2^2 \\
 \text{s.t.} & x_1^2 + x_2^2 \leq 25 \\
 & x_1 \geq 0
 \end{array}$$

Exercise 5 (When KKT can fail without qualification). Consider the problem

$$\min_{x \in \mathbb{R}} f(x) := x \quad \text{s.t.} \quad g(x) := x^2 \leq 0.$$

1. Find the (global) solution.
2. Write the KKT conditions at the solution and show that there is no Lagrange multiplier $\mu \geq 0$ satisfying them.
3. Compute the tangent cone $T_X(x^*)$ and the linearized cone $T_X^\ell(x^*)$ at the solution x^* (with $X = \{x : g(x) \leq 0\}$).