

## Exercises: Constrained Optimization

**Exercise 1** (Penalization). We consider the following problem

$$(P) \quad \begin{aligned} \min_{x \in \mathbb{R}^n} \quad & f(x) \\ \text{s.t.} \quad & Ax = b, \quad x \leq 0 \end{aligned}$$

with value  $v$  and the following penalized versions

$$(P_t^{in}) \quad \begin{aligned} \min_{x \in \mathbb{R}^n} \quad & f(x) - t \sum_{i=1}^n \ln(-x_i) \\ \text{s.t.} \quad & Ax = b, \quad x < 0 \end{aligned}$$

and

$$(P_t^{out}) \quad \begin{aligned} \min_{x \in \mathbb{R}^n} \quad & f(x) + t \sum_{i=1}^n (x_i)^+ \\ \text{s.t.} \quad & Ax = b \end{aligned}$$

with associated value  $v_t^{in}$  and  $v_t^{out}$ , and an optimal solution  $x_t^{in}$  and  $x_t^{out}$ .

1. Intuitively, assuming that  $f$  is "well behaved", for  $t$  going to which value does  $(P_t^{in})$  tend to the original problem  $(P)$ ? In which sense?
2. What can you say about  $x_t^{in}$ ?
3. Can you compare  $v_t^{in}$  and  $v$ ?
4. Same questions for  $(P_t^{out})$ .

**Exercise 2** (Decomposition by prices). We consider the following energy problem:

- you are an energy producer with  $N$  production units
- you have to satisfy a given demand planning for the next 24h (i.e. the total output at time  $t$  should be equal to  $d_t$ )

- the time step is the hour, and each unit have a production cost for each planning given as a convex quadratic function of the planning
- For each unit  $i$ , the production planning  $u^i = (u_t^i)_{t \in [24]}$  has to satisfy polyhedral constraints  $u^i \in U^i$ .

1. Model this problem as an optimization problem. In which class does it belongs? How many variables?
2. Apply Uzawa's algorithm to this problem. Why could this be an interesting idea?
3. Give an economic interpretation to this method.
4. What would happen if each unit had production constraints?

**Exercise 3** (Kelley's convergence). We are going to prove that, if  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is convex, and  $X$  a non-empty polytope (bounded polyhedron) then Kelley's cutting plane algorithm is converging.

Consider  $x_1 \in X$ . We consider a sequence of points  $(x^{(k)})_{k \in \mathbb{N}}$  such that  $x^{(k+1)}$  is an optimal solution to

$$(P^{(k)}) \quad \begin{aligned} \underline{v}^{(k+1)} = \text{Min}_{x \in X} \quad & z \\ \text{s.t.} \quad & f(x^{(\kappa)}) + \langle g^{(\kappa)}, x - x^{(\kappa)} \rangle \leq z \quad \forall \kappa \in [k] \end{aligned}$$

where  $g^{(k)} \in \partial f(x^{(k)})$ .

Denote  $v = \min_{x \in X} f(x)$ .

1. Show that  $v$  exists and is finite, and that there exists a sequence  $x^{(k)}$ .
2. Show that there exists  $L$  such that, for all  $k_1$  and  $k_2$ , we have  $\|f(x^{(k_1)}) - f(x^{(k_2)})\| \leq L\|x^{(k_1)} - x^{(k_2)}\|$ , and  $\|g^{(k)}\| \leq L$ .

3. Let  $K_\varepsilon = \{k \in \mathbb{N} \mid f(x^{(k)}) > v + \varepsilon\}$  be the set of index such that  $x^{(k)}$  is not an  $\varepsilon$ -optimal solution. Show that  $f(x_k) \rightarrow v$  if and only if  $K_\varepsilon$  is finite for all  $\varepsilon > 0$
4. Consider  $k_1, k_2 \in K_\varepsilon$ , such that  $k_2 > k_1$ . Show that
- $$f(x^{(k_1)}) + \langle g^{(k_1)}, x^{(k_2)} - x^{(k_1)} \rangle \leq \underline{v}^{(k_2)} \leq v$$
5. Show that  $\varepsilon + f(x^{(k_1)}) + \langle g^{(k_1)}, x^{(k_2)} - x^{(k_1)} \rangle < f(x^{(k_2)})$
6. Show that  $\varepsilon < 2L\|x^{(k_2)} - x^{(k_1)}\|$ .
7. Prove that  $f(x^{(k)}) \rightarrow v$ .
8. (Optional - hard) Find a complexity bound for the method (that is a number of iteration  $N_\varepsilon$  after which you are sure to have obtained a  $\varepsilon$ -optimal solution).